

**Systematic Review of Management Options for Atrial Fibrillation
Citation List**

Catheter ablation

1. Arentz T, Jander N, von Rosenthal J, et al. Incidence of pulmonary vein stenosis 2 years after radiofrequency catheter ablation of refractory atrial fibrillation. *European Heart Journal*. 2003;24(10):963-969.
2. Arentz T, von Rosenthal J, Blum T, et al. Feasibility and safety of pulmonary vein isolation using a new mapping and navigation system in patients with refractory atrial fibrillation. *Circulation*. 2003;108(20):2484-2490.
3. Atienza F, Almendral J, Jalife J, et al. Real-time dominant frequency mapping and ablation of dominant frequency sites in atrial fibrillation with left-to-right frequency gradients predicts long-term maintenance of sinus rhythm. *Heart Rhythm*. 2009;6(1):33-40.
4. Baman TS, Gupta SK, Billakanty SR, et al. Time to cardioversion of recurrent atrial arrhythmias after catheter ablation of atrial fibrillation and long-term clinical outcome. *J Cardiovasc Electrophysiol*. Dec 2009;20(12):1321-1325.
5. Berkowitsch A, Neumann T, Kuniss M, Brandt R, Zaltberg S, Pitschner HF. Evaluation of atrial fibrillation burden before catheter ablation predicts outcome after pulmonary vein isolation. *Indian Pacing Electrophysiol. J*. 2009;9(3):138-150.
6. Bertaglia E, Bella PD, Tondo C, et al. Image integration increases efficacy of paroxysmal atrial fibrillation catheter ablation: results from the CartoMerge Italian Registry. *Europace: European Pacing, Arrhythmias, and Cardiac Electrophysiology: Journal of the Working Groups on Cardiac Pacing, Arrhythmias, and Cardiac Cellular Electrophysiology of the European Society of Cardiology*. 2009;11(8):1004-1010.
7. Bunch TJ, Weiss JP, Crandall BG, et al. Long-Term Clinical Efficacy and Risk of Catheter Ablation for Atrial Fibrillation in Octogenarians. *Pacing and Clinical Electrophysiology: PACE*. 2009.

8. Calvo N, Mont L, Tamborero D, et al. Efficacy of circumferential pulmonary vein ablation of atrial fibrillation in endurance athletes. *Europace: European Pacing, Arrhythmias, and Cardiac Electrophysiology: Journal of the Working Groups on Cardiac Pacing, Arrhythmias, and Cardiac Cellular Electrophysiology of the European Society of Cardiology*. 2010;12(1):30-36.
9. Cheema A, Dong J, Dalal D, et al. Long-term safety and efficacy of circumferential ablation with pulmonary vein isolation. *Journal of cardiovascular electrophysiology*. 2006;17(10):1080-1085.
10. Corrado A, Bonso A, Madalosso M, et al. Impact of systematic isolation of superior vena cava in addition to pulmonary vein antrum isolation on the outcome of paroxysmal, persistent, and permanent atrial fibrillation ablation: Results from a randomized study. *J. Cardiovasc. Electrophysiol*. 2010;21(1):1-5.
11. Deisenhofer I, Estner H, Reents T, et al. Does electrogram guided substrate ablation add to the success of pulmonary vein isolation in patients with paroxysmal atrial fibrillation? A prospective, randomized study. *Journal of cardiovascular electrophysiology*. 2009;20(5):514-521.
12. Deisenhofer I, Schneider MAE, Böhlen-Knauf M, et al. Circumferential mapping and electric isolation of pulmonary veins in patients with atrial fibrillation. *The American journal of cardiology*. 2003;91(2):159-163.
13. Della Bella P, Fassini G, Cireddu M, et al. Image integration-guided catheter ablation of atrial fibrillation: a prospective randomized study. *Journal of cardiovascular electrophysiology*. 2009;20(3):258-265.
14. Di Biase L, Elayi CS, Fahmy TS, et al. Atrial fibrillation ablation strategies for paroxysmal patients: randomized comparison between different techniques. *Circulation. Arrhythmia and electrophysiology*. 2009;2(2):113-119.
15. Di Biase L, Wang Y, Horton R, et al. Ablation of atrial fibrillation utilizing robotic catheter navigation in comparison to manual navigation and ablation: Single-center experience. *J. Cardiovasc. Electrophysiol*. 2009;20(12):1328-1335.
16. Essebag V, Wylie JV, Reynolds MR, et al. Bi-directional electrical pulmonary vein isolation as an endpoint for ablation of paroxysmal atrial fibrillation. *Journal of Interventional Cardiac Electrophysiology: An International Journal of Arrhythmias and Pacing*. 2006;17(2):111-117.
17. Forleo GB, Mantica M, De Luca L, et al. Catheter ablation of atrial fibrillation in patients with diabetes mellitus type 2: results from a randomized study comparing pulmonary vein isolation versus antiarrhythmic drug therapy. *Journal of cardiovascular electrophysiology*. 2009;20(1):22-28.
18. Haïssaguerre M, Jaïs P, Shah DC, et al. Spontaneous initiation of atrial fibrillation by ectopic beats originating in the pulmonary veins. *The New England journal of medicine*. 1998;339(10):659-666.

19. Helms AS, West JJ, Patel A, et al. Relation of left atrial volume from three-dimensional computed tomography to atrial fibrillation recurrence following ablation. *The American journal of cardiology*. 2009;103(7):989-993.
20. Hocini M, Jaïs P, Sanders P, et al. Techniques, evaluation, and consequences of linear block at the left atrial roof in paroxysmal atrial fibrillation: a prospective randomized study. *Circulation*. 2005;112(24):3688-3696.
21. Hof I, Chilukuri K, Arbab-Zadeh A, et al. Does left atrial volume and pulmonary venous anatomy predict the outcome of catheter ablation of atrial fibrillation? *Journal of cardiovascular electrophysiology*. 2009;20(9):1005-1010.
22. Hsieh M-H, Tai C-T, Lee S-H, et al. Catheter ablation of atrial fibrillation versus atrioventricular junction ablation plus pacing therapy for elderly patients with medically refractory paroxysmal atrial fibrillation. *Journal of cardiovascular electrophysiology*. 2005;16(5):457-461.
23. Husser D, Bollmann A, Kang S, et al. Effectiveness of catheter ablation for coexisting atrial fibrillation and atrial flutter. *The American journal of cardiology*. 2004;94(5):666-668.
24. Jaïs P, Cauchemez B, Macle L, et al. Catheter ablation versus antiarrhythmic drugs for atrial fibrillation: the A4 study. *Circulation*. 2008;118(24):2498-2505.
25. Joshi S, Choi AD, Kamath GS, et al. Prevalence, predictors, and prognosis of atrial fibrillation early after pulmonary vein isolation: findings from 3 months of continuous automatic ECG loop recordings. *Journal of cardiovascular electrophysiology*. 2009;20(10):1089-1094.
26. Kanagaratnam L, Tomassoni G, Schweikert R, et al. Empirical pulmonary vein isolation in patients with chronic atrial fibrillation using a three-dimensional nonfluoroscopic mapping system: long-term follow-up. *Pacing and Clinical Electrophysiology: PACE*. 2001;24(12):1774-1779.
27. Khaykin Y, Skanes A, Champagne J, et al. A randomized controlled trial of the efficacy and safety of electroanatomic circumferential pulmonary vein ablation supplemented by ablation of complex fractionated atrial electrograms versus potential-guided pulmonary vein antrum isolation guided by intracardiac ultrasound. *Circulation. Arrhythmia and electrophysiology*. 2009;2(5):481-487.
28. Klemm HU, Ventura R, Rostock T, et al. Correlation of symptoms to ECG diagnosis following atrial fibrillation ablation. *Journal of cardiovascular electrophysiology*. 2006;17(2):146-150.
29. Kriatselis C, Tang M, Nedios S, Roser M, Gerds-Li H, Fleck E. Intraprocedural reconstruction of the left atrium and pulmonary veins as a single navigation tool for ablation of atrial fibrillation: a feasibility, efficacy, and safety study. *Heart Rhythm: The Official Journal of the Heart Rhythm Society*. 2009;6(6):733-741.

30. Kumagai K, Ogawa M, Noguchi H, et al. Comparison of 2 mapping strategies for pulmonary vein isolation. *Circulation Journal: Official Journal of the Japanese Circulation Society*. 2005;69(12):1496-1502.
31. Kusumoto F, Prussak K, Wiesinger M, Pullen T, Lynady C. Radiofrequency catheter ablation of atrial fibrillation in older patients: outcomes and complications. *Journal of Interventional Cardiac Electrophysiology: An International Journal of Arrhythmias and Pacing*. 2009;25(1):31-35.
32. Lin Y-J, Tai C-T, Chang S-L, et al. Efficacy of additional ablation of complex fractionated atrial electrograms for catheter ablation of nonparoxysmal atrial fibrillation. *Journal of cardiovascular electrophysiology*. 2009;20(6):607-615.
33. Lo LW, Lin YJ, Tsao HM, et al. The impact of left atrial size on long-term outcome of catheter ablation of chronic atrial fibrillation. *J. Cardiovasc. Electrophysiol*. 2009;20(11):1211-1216.
34. Macle L, Jaïs P, Weerasooriya R, et al. Irrigated-tip catheter ablation of pulmonary veins for treatment of atrial fibrillation. *Journal of cardiovascular electrophysiology*. 2002;13(11):1067-1073.
35. Macle L, Novak P, Khairy P, et al. Pulmonary vein isolation and left atrial catheter ablation using a three-dimensional navigation system for the treatment of atrial fibrillation. *The Canadian Journal of Cardiology*. 2009;25(6):e169-172-e169-172.
36. Malmborg H, Lönnerholm S, Blomström-Lundqvist C. Acute and clinical effects of cryoballoon pulmonary vein isolation in patients with symptomatic paroxysmal and persistent atrial fibrillation. *Europace: European Pacing, Arrhythmias, and Cardiac Electrophysiology: Journal of the Working Groups on Cardiac Pacing, Arrhythmias, and Cardiac Cellular Electrophysiology of the European Society of Cardiology*. 2008;10(11):1277-1280.
37. Mangrum JM, Mounsey JP, Kok LC, DiMarco JP, Haines DE. Intracardiac echocardiography-guided, anatomically based radiofrequency ablation of focal atrial fibrillation originating from pulmonary veins. *Journal of the American College of Cardiology*. 2002;39(12):1964-1972.
38. Marrouche NF, Guenther J, Segerson NM, et al. Randomized comparison between open irrigation technology and intracardiac-echo-guided energy delivery for pulmonary vein antrum isolation: procedural parameters, outcomes, and the effect on esophageal injury. *Journal of cardiovascular electrophysiology*. 2007;18(6):583-588.
39. Matsuo S, Lellouche N, Wright M, et al. Clinical predictors of termination and clinical outcome of catheter ablation for persistent atrial fibrillation. *Journal of the American College of Cardiology*. 2009;54(9):788-795.

40. Meissner A, van Bracht M, Schrage M-O, et al. Segmental pulmonary vein isolation in atrial fibrillation: new insights from the high density mesh mapper technique in an electrophysiologically guided approach. *Journal of Interventional Cardiac Electrophysiology: An International Journal of Arrhythmias and Pacing*. 2009;25(3):183-192.
41. Mesas CE, Augello G, Lang CCE, et al. Electroanatomic remodeling of the left atrium in patients undergoing repeat pulmonary vein ablation: mechanistic insights and implications for ablation. *Journal of cardiovascular electrophysiology*. 2006;17(12):1279-1285.
42. Neumann T, Vogt J, Schumacher B, et al. Circumferential pulmonary vein isolation with the cryoballoon technique results from a prospective 3-center study. *Journal of the American College of Cardiology*. 2008;52(4):273-278.
43. O'Neill MD, Wright M, Knecht S, et al. Long-term follow-up of persistent atrial fibrillation ablation using termination as a procedural endpoint. *European Heart Journal*. 2009;30(9):1105-1112.
44. Oral H, Pappone C, Chugh A, et al. Circumferential pulmonary-vein ablation for chronic atrial fibrillation. *N Engl J Med*. Mar 2 2006;354(9):934-941.
45. Pappone C, Augello G, Sala S, et al. A randomized trial of circumferential pulmonary vein ablation versus antiarrhythmic drug therapy in paroxysmal atrial fibrillation: the APAF Study. *Journal of the American College of Cardiology*. 2006;48(11):2340-2347.
46. Pappone C, Oreto G, Rosanio S, et al. Atrial electroanatomic remodeling after circumferential radiofrequency pulmonary vein ablation: efficacy of an anatomic approach in a large cohort of patients with atrial fibrillation. *Circulation*. 2001;104(21):2539-2544.
47. Pappone C, Rosanio S, Augello G, et al. Mortality, morbidity, and quality of life after circumferential pulmonary vein ablation for atrial fibrillation: outcomes from a controlled nonrandomized long-term study. *J Am Coll Cardiol*. Jul 16 2003;42(2):185-197.
48. Patel D, Mohanty P, Di Biase L, et al. Outcomes and complications of catheter ablation for atrial fibrillation in females. *Heart Rhythm: The Official Journal of the Heart Rhythm Society*. 2010;7(2):167-172.
49. Pontoppidan J, Nielsen JC, Poulsen SH, et al. Prophylactic cavotricuspid isthmus block during atrial fibrillation ablation in patients without atrial flutter: a randomised controlled trial. *Heart (British Cardiac Society)*. 2009;95(12):994-999.
50. Rossillo A, Bonso A, Themistoclakis S, et al. Role of anticoagulation therapy after pulmonary vein antrum isolation for atrial fibrillation treatment. *J Cardiovasc Med (Hagerstown)*. Jan 2008;9(1):51-55.

51. Saad EB, Rossillo A, Saad CP, et al. Pulmonary vein stenosis after radiofrequency ablation of atrial fibrillation: functional characterization, evolution, and influence of the ablation strategy. *Circulation*. 2003;108(25):3102-3107.
52. Saliba W, Reddy VY, Wazni O, et al. Atrial fibrillation ablation using a robotic catheter remote control system: initial human experience and long-term follow-up results. *Journal of the American College of Cardiology*. 2008;51(25):2407-2411.
53. Sawhney N, Anousheh R, Chen W-C, Narayan S, Feld GK. Five-year outcomes after segmental pulmonary vein isolation for paroxysmal atrial fibrillation. *The American journal of cardiology*. 2009;104(3):366-372.
54. Scharf C, Boersma L, Davies W, et al. Ablation of persistent atrial fibrillation using multielectrode catheters and duty-cycled radiofrequency energy. *Journal of the American College of Cardiology*. 2009;54(15):1450-1456.
55. Shin S-H, Park M-Y, Oh W-J, et al. Left atrial volume is a predictor of atrial fibrillation recurrence after catheter ablation. *Journal of the American Society of Echocardiography: Official Publication of the American Society of Echocardiography*. 2008;21(6):697-702.
56. Siklody CH, Minners J, Allgeier M, et al. Cryoballoon pulmonary vein isolation guided by transesophageal echocardiography: Novel aspects on an emerging ablation technique. *J. Cardiovasc. Electrophysiol*. 2009;20(11):1197-1202.
57. Stabile G, Bertaglia E, Senatore G, et al. Catheter ablation treatment in patients with drug-refractory atrial fibrillation: a prospective, multi-centre, randomized, controlled study (Catheter Ablation For The Cure Of Atrial Fibrillation Study). *Eur Heart J*. Jan 2006;27(2):216-221.
58. Stabile G, Bertaglia E, Turco P, et al. Role of pulmonary veins isolation in persistent atrial fibrillation ablation: the pulmonary vein isolation in persistent atrial fibrillation (PIPA) study. *Pacing and Clinical Electrophysiology: PACE*. 2009;32 Suppl 1:S116-119-S116-119.
59. Steven D, Servatius H, Rostock T, et al. Reduced fluoroscopy during atrial fibrillation ablation: Benefits of robotic guided navigation. *J. Cardiovasc. Electrophysiol*. 2010;21(1):6-12.
60. Tamborero D, Mont L, Berruezo A, et al. Left atrial posterior wall isolation does not improve the outcome of circumferential pulmonary vein ablation for atrial fibrillation: a prospective randomized study. *Circulation. Arrhythmia and electrophysiology*. 2009;2(1):35-40.
61. Tan ES, Mulder BA, Rienstra M, et al. Pulmonary vein isolation of symptomatic refractory paroxysmal and persistent atrial fibrillation: A single centre and single operator experience in the Netherlands. *Neth. Heart J*. 2009;17(10):366-372.

62. Udyavar AR, Chang S-L, Tai C-T, et al. The important role of pulmonary vein carina ablation as an adjunct to circumferential pulmonary vein isolation. *Journal of cardiovascular electrophysiology*. 2008;19(6):593-598.
63. Van Belle Y, Janse P, Theuns D, Szili-Torok T, Jordaens L. One year follow-up after cryoballoon isolation of the pulmonary veins in patients with paroxysmal atrial fibrillation. *Europace: European Pacing, Arrhythmias, and Cardiac Electrophysiology: Journal of the Working Groups on Cardiac Pacing, Arrhythmias, and Cardiac Cellular Electrophysiology of the European Society of Cardiology*. 2008;10(11):1271-1276.
64. Wazni OM, Barrett C, Martin DO, et al. Experience with the Hansen robotic system for atrial fibrillation ablation - Lessons learned and techniques modified: Hansen in the real world. *J. Cardiovasc. Electrophysiol*. 2009;20(11):1193-1196.
65. Wazni OM, Marrouche NF, Martin DO, et al. Radiofrequency ablation vs antiarrhythmic drugs as first-line treatment of symptomatic atrial fibrillation: a randomized trial. *JAMA*. Jun 1 2005;293(21):2634-2640.
66. Wiesfeld ACP, Tan ES, Van Veldhuisen DJ, Crijns HJGM, Van Gelder IC. Radiofrequency ablation of atrial fibrillation. *International Journal of Cardiology*. 2004;93(2-3):231-237.
67. Wilber DJ, Pappone C, Neuzil P, et al. Comparison of antiarrhythmic drug therapy and radiofrequency catheter ablation in patients with paroxysmal atrial fibrillation: A randomized controlled trial. *J. Am. Med. Assoc*. 2010;303(4):333-340.
68. Yamada T, Murakami Y, Okada T, et al. Electrophysiological pulmonary vein antrum isolation with a multielectrode basket catheter is feasible and effective for curing paroxysmal atrial fibrillation: efficacy of minimally extensive pulmonary vein isolation. *Heart Rhythm: The Official Journal of the Heart Rhythm Society*. 2006;3(4):377-384.
69. Yamada T, Yoshida N, Murakami Y, et al. Vagal modification can be a valid predictor of late recurrence of paroxysmal atrial fibrillation independent of the pulmonary vein isolation technique. *Circulation Journal: Official Journal of the Japanese Circulation Society*. 2009;73(9):1606-1611.
70. Yoshida N, Yamada T, Murakami Y, et al. Vagal modification can also help prevent late recurrence of atrial fibrillation after segmental pulmonary vein isolation. *Circulation Journal: Official Journal of the Japanese Circulation Society*. 2009;73(4):632-638.

Antiarrhythmic agents

1. Maintenance of sinus rhythm in patients with atrial fibrillation: an AFFIRM substudy of the first antiarrhythmic drug. *Journal of the American College of Cardiology*. 2003;42(1):20-29.
2. Aizawa Y, Kohsaka S, Suzuki S, et al. Comparison of antiarrhythmics used in patients with paroxysmal atrial fibrillation. *Circulation Journal: Official Journal of the Japanese Circulation Society*. 2010;74(1):71-76.
3. Banchs JE, Wolbrette DL, Samii SM, et al. Efficacy and safety of dofetilide in patients with atrial fibrillation and atrial flutter. *Journal of Interventional Cardiac Electrophysiology: An International Journal of Arrhythmias and Pacing*. 2008;23(2):111-115.
4. Bellandi F, Simonetti I, Leoncini M, et al. Long-term efficacy and safety of propafenone and sotalol for the maintenance of sinus rhythm after conversion of recurrent symptomatic atrial fibrillation. *The American journal of cardiology*. 2001;88(6):640-645.
5. Carlsson J, Miketic S, Windeler J, et al. Randomized trial of rate-control versus rhythm-control in persistent atrial fibrillation: the Strategies of Treatment of Atrial Fibrillation (STAF) study. *Journal of the American College of Cardiology*. 2003;41(10):1690-1696.
6. Channer KS, Birchall A, Steeds RP, et al. A randomized placebo-controlled trial of pre-treatment and short- or long-term maintenance therapy with amiodarone supporting DC cardioversion for persistent atrial fibrillation. *European Heart Journal*. 2004;25(2):144-150.
7. Connolly SJ, Crijns HJGM, Torp-Pedersen C, et al. Analysis of stroke in ATHENA: a placebo-controlled, double-blind, parallel-arm trial to assess the efficacy of dronedarone 400 mg BID for the prevention of cardiovascular hospitalization or death from any cause in patients with atrial fibrillation/atrial flutter. *Circulation*. 2009;120(13):1174-1180.
8. Hohnloser SH, Crijns HJGM, van Eickels M, et al. Effect of dronedarone on cardiovascular events in atrial fibrillation. *The New England journal of medicine*. 2009;360(7):668-678.
9. Hohnloser SH, Kuck KH, Lilienthal J. Rhythm or rate control in atrial fibrillation-Pharmacological Intervention in Atrial Fibrillation (PIAF): a randomised trial. *Lancet*. 2000;356(9244):1789-1794.
10. Kilborn MJ, Rathore SS, Gersh BJ, Oetgen WJ, Solomon AJ. Amiodarone and mortality among elderly patients with acute myocardial infarction with atrial fibrillation. *American heart journal*. 2002;144(6):1095-1101.
11. Kochiadakis GE, Igoumenidis NE, Hamilos ME, et al. Sotalol versus propafenone for long-term maintenance of normal sinus rhythm in patients with recurrent

- symptomatic atrial fibrillation. *The American journal of cardiology*. 2004;94(12):1563-1566.
12. Kochiadakis GE, Igoumenidis NE, Hamilos MI, et al. Long-term maintenance of normal sinus rhythm in patients with current symptomatic atrial fibrillation: amiodarone vs propafenone, both in low doses. *Chest*. 2004;125(2):377-383.
 13. Kochiadakis GE, Igoumenidis NE, Marketou ME, Kaleboubas MD, Simantirakis EN, Vardas PE. Low dose amiodarone and sotalol in the treatment of recurrent, symptomatic atrial fibrillation: a comparative, placebo controlled study. *Heart (British Cardiac Society)*. 2000;84(3):251-257.
 14. Li H, Riedel R, Oldemeyer JB, Rovang K, Hee T. Comparison of recurrence rates after direct-current cardioversion for new-onset atrial fibrillation in patients receiving versus those not receiving rhythm-control drug therapy. *The American journal of cardiology*. 2004;93(1):45-48.
 15. Nergårdh A, Nordlander R, Frick M. Rate of conversion and recurrence after sotalol treatment in patients with direct current-refractory atrial fibrillation. *Clinical cardiology*. 2006;29(2):56-60.
 16. Ogawa S, Yamashita T, Yamazaki T, et al. Optimal treatment strategy for patients with paroxysmal atrial fibrillation: J-RHYTHM Study. *Circulation Journal: Official Journal of the Japanese Circulation Society*. 2009;73(2):242-248.
 17. Plewan A, Lehmann G, Ndrepepa G, et al. Maintenance of sinus rhythm after electrical cardioversion of persistent atrial fibrillation; sotalol vs bisoprolol. *European Heart Journal*. 2001;22(16):1504-1510.
 18. Pritchett ELC, Page RL, Carlson M, Undesser K, Fava G. Efficacy and safety of sustained-release propafenone (propafenone SR) for patients with atrial fibrillation. *The American journal of cardiology*. 2003;92(8):941-946.
 19. Roy D, Talajic M, Dorian P, et al. Amiodarone to prevent recurrence of atrial fibrillation. Canadian Trial of Atrial Fibrillation Investigators. *The New England journal of medicine*. 2000;342(13):913-920.
 20. Roy D, Talajic M, Nattel S, et al. Rhythm control versus rate control for atrial fibrillation and heart failure. *The New England journal of medicine*. 2008;358(25):2667-2677.
 21. Sanofi-Aventis. *Multaq (dronedarone) briefing document for the advisory committee meeting of the Cardiovascular and Renal Drugs Division of the US Food and Drug Administration* March 18, 2009 2009.
 22. Singh BN, Connolly SJ, Crijns HJGM, et al. Dronedarone for maintenance of sinus rhythm in atrial fibrillation or flutter. *The New England journal of medicine*. 2007;357(10):987-999.

23. Singh BN, Singh SN, Reda DJ, et al. Amiodarone versus sotalol for atrial fibrillation. *The New England journal of medicine*. 2005;352(18):1861-1872.
24. Singh S, Zoble RG, Yellen L, et al. Efficacy and safety of oral dofetilide in converting to and maintaining sinus rhythm in patients with chronic atrial fibrillation or atrial flutter: the symptomatic atrial fibrillation investigative research on dofetilide (SAFIRE-D) study. *Circulation*. 2000;102(19):2385-2390.
25. Touboul P, Brugada J, Capucci A, Crijns HJGM, Edvardsson N, Hohnloser SH. Dronedarone for prevention of atrial fibrillation: a dose-ranging study. *European Heart Journal*. 2003;24(16):1481-1487.
26. Tse H-F, Lau CP. Clinical predictors and time course of arrhythmia recurrence in patients with early reinitiation of atrial fibrillation after successful internal cardioversion. *Pacing and Clinical Electrophysiology: PACE*. 2003;26(9):1809-1814.
27. Van Gelder IC, Hagens VE, Bosker HA, et al. A comparison of rate control and rhythm control in patients with recurrent persistent atrial fibrillation. *The New England journal of medicine*. 2002;347(23):1834-1840.

Surgical ablation

1. Akpınar B, Guden M, Sagbas E, et al. Combined radiofrequency modified maze and mitral valve procedure through a port access approach: early and mid-term results. *European Journal of Cardio-Thoracic Surgery: Official Journal of the European Association for Cardio-Thoracic Surgery*. 2003;24(2):223-230.
2. Beyer E, Lee R, Lam B-K. Point: Minimally invasive bipolar radiofrequency ablation of lone atrial fibrillation: early multicenter results. *The Journal of Thoracic and Cardiovascular Surgery*. 2009;137(3):521-526.
3. Cui YQ, Sun LB, Li Y, et al. Intraoperative modified Cox mini-maze procedure for long-standing persistent atrial fibrillation. *The Annals of Thoracic Surgery*. 2008;85(4):1283-1289.
4. Edgerton JR, Jackman WM, Mahoney C, Mack MJ. Totally thorascopic surgical ablation of persistent AF and long-standing persistent atrial fibrillation using the "Dallas" lesion set. *Heart Rhythm*. 2009;6(12 SUPPL.):S64-S70-S64-S70.
5. Edgerton JR, McClelland JH, Duke D, et al. Minimally invasive surgical ablation of atrial fibrillation: six-month results. *The Journal of Thoracic and Cardiovascular Surgery*. 2009;138(1):109-113; discussion 114-109-113; discussion 114.
6. Geidel S, Lass M, Ostermeyer J. A 5-year clinical experience with bipolar radiofrequency ablation for permanent atrial fibrillation concomitant to coronary artery bypass grafting and aortic valve surgery. *Interactive Cardiovascular and Thoracic Surgery*. 2008;7(5):777-780.
7. Han FT, Kasirajan V, Kowalski M, et al. Results of a minimally invasive surgical pulmonary vein isolation and ganglionic plexi ablation for atrial fibrillation: single-center experience with 12-month follow-up. *Circulation. Arrhythmia and electrophysiology*. 2009;2(4):370-377.
8. Onorati F, Esposito A, Messina G, di Virgilio A, Renzulli A. Right isthmus ablation reduces supraventricular arrhythmias after surgery for chronic atrial fibrillation. *The Annals of Thoracic Surgery*. 2008;85(1):39-48.
9. Wolf RK, Schneeberger EW, Osterday R, et al. Video-assisted bilateral pulmonary vein isolation and left atrial appendage exclusion for atrial fibrillation. *The Journal of Thoracic and Cardiovascular Surgery*. 2005;130(3):797-802.
10. Yilmaz A, Geuzebroek GSC, Van Putte BP, et al. Completely thorascopic pulmonary vein isolation with ganglionic plexus ablation and left atrial appendage amputation for treatment of atrial fibrillation. *European Journal of Cardio-Thoracic Surgery: Official Journal of the European Association for Cardio-Thoracic Surgery*. 2010.